

1.1 – Introduction to Systems of Linear Equations

A **linear equation** can be expressed in the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$$

where a_i are constants, not all zero, and b is a constant.

14. In each part, use parametric equations to describe the solution set of the linear equation.

a. $x + 10y = 2$

b. $x_1 + 3x_2 - 12x_3 = 3$

c. $4x_1 + 2x_2 + 3x_3 + x_4 = 20$

d. $v + w + x - 5y + 7z = 0$

a. $x = -10y + 2$

Let $t = y$.

Then

$$x = -10t + 2$$

$$y = t$$

c. $x_1 = -\frac{1}{2}x_2 - \frac{3}{4}x_3 - \frac{1}{4}x_4 + 5$

Let $r = x_2, s = x_3, t = x_4$

Then

$$x_1 = -\frac{1}{2}r - \frac{3}{4}s - \frac{1}{4}t + 5$$

$$x_2 = r$$

$$x_3 = s$$

$$x_4 = t$$

A **homogeneous linear equation** is a linear equation in which $b = 0$.

A **system of linear equations** is a finite set of linear equations. It has the form

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2 \\ \vdots & \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= b_m \end{aligned}$$

Note that the first of the double subscripts references the equation number, and the second references the term in that equation.

The **augmented matrix** for the above linear system is

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \vdots & \vdots & & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \vdots & \vdots & & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{bmatrix}$$

16. Each linear system has infinitely many solutions. Use parametric equations to describe its solution set.

a.

$$6x_1 + 2x_2 = -8$$

$$3x_1 + x_2 = -4$$

$$\text{EQ 1} \rightarrow \frac{1}{2} \text{EQ 1} \quad 3x_1 + x_2 = -4$$

$$3x_1 + x_2 = -4$$

$$x_1 = -\frac{1}{3}x_2 - \frac{4}{3}$$

$$x_1 = -\frac{1}{3}t - \frac{4}{3}$$

$$x_2 = t$$

Augmented matrix

$$\left[\begin{array}{cc|c} 6 & 2 & -8 \\ 3 & 1 & -4 \end{array} \right]$$

$$R_1 \rightarrow \frac{1}{2}R_1$$

$$\left[\begin{array}{cc|c} 3 & 1 & -4 \\ 3 & 1 & -4 \end{array} \right] \quad R_2 \rightarrow R_2 - R_1$$

$$\left[\begin{array}{cc|c} 3 & 1 & -4 \\ 0 & 0 & 0 \end{array} \right]$$

b.

$$\begin{array}{l} 2x - y + 2z = -4 \\ 6x - 3y + 6z = -12 \\ -4x + 2y - 4z = 8 \end{array} \rightarrow \begin{array}{l} 2x - y + 2z = -4 \\ 2x - y + 2z = -4 \\ 2x - y + 2z = -4 \end{array}$$

$$x = \frac{1}{2}y - z - 2 \quad \text{Let } s = y \text{ and } t = z.$$

$$\begin{array}{l} x = \frac{1}{2}s - t - 2 \\ y = s \\ z = t \end{array}$$

Alternative form: $(x, y, z) = (\frac{1}{2}s - t - 2, s, t)$

The following are **elementary row operations** performed on a matrix.

1. Multiply a row by a nonzero constant.
2. Interchange two rows.
3. Add a constant times one row to another.

A **solution** of a linear system is a sequence of n numbers s_1, s_2, \dots, s_n that when substituted for corresponding unknowns x_i makes each equation a true statement. If $n = 2$, then the solution is an **ordered pair**, and if $n = 3$, it is an **ordered triple**. In general, an **ordered n -tuple** has the form (s_1, s_2, \dots, s_n) .

A linear system is **consistent** if it has at least one solution.

A linear system is **inconsistent** if it has no solutions.

20. Find all values of k for which the given augmented matrix corresponds to a consistent linear system.

a. $\left[\begin{array}{cc|c} 3 & -4 & k \\ -6 & 8 & 5 \end{array} \right]$

$$R_2 \rightarrow R_2 + 2R_1$$

$$\begin{array}{cc|c} -6 & 8 & 5 \\ 6 & -8 & 2k \\ \hline 0 & 0 & 2k+5 \end{array}$$

$$\left[\begin{array}{cc|c} \overset{x}{3} & \overset{y}{-4} & k \\ 0 & 0 & 2k+5 \end{array} \right]$$

$$0 = 2k+5$$

$$k = -\frac{5}{2}$$

b. $\left[\begin{array}{cc|c} \overset{x}{k} & \overset{y}{1} & -2 \\ 4 & -1 & 2 \end{array} \right] \rightarrow 4x - y = 2$

$$R_2 \rightarrow R_2 + R_1$$

$$\left[\begin{array}{cc|c} k & 1 & -2 \\ k+4 & 0 & 0 \end{array} \right]$$

$$(k+4)x = 0$$

$$\downarrow \quad \rightarrow \quad x = 0$$
$$k = -4$$

k can be anything

k can be any number.

If $x=0$,
 $y=-2$

Ex: A 3-7-9 diet calls for 3 units of fat, 7 units of protein, and 9 units of carbs in each meal. Suppose an individual has three possible foods to choose from to meet these requirements. Each ounce of the food contains

x	Food 1: 3 units of fat, 4 units of protein, and 1 unit of carbs
y	Food 2: 2 units of fat, 5 units of protein, 3 units of carbs
z	Food 3: 4 units of fat, 1 unit of protein, 2 units of carbs

Let x , y , and z denote the number of ounces of the first, second, and third foods that a person will consume at the main meal. Find a linear system in x , y , and z whose solution tells how many ounces of each food must be consumed to meet the diet requirements.

$$\begin{array}{l} \text{Fat:} \\ \text{protein:} \\ \text{Carbs:} \end{array} \quad \begin{array}{l} 3x + 2y + 4z = 3 \\ 4x + 5y + z = 7 \\ x + 3y + 2z = 9 \end{array}$$