

4.3 – Spanning Sets

Due Sun

Definition: (generalizes linear combination from Section 3.1)

If \mathbf{w} is a vector in a vector space V , then \mathbf{w} is said to be a **linear combination** of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r$ in V if \mathbf{w} can be expressed in the form

$\mathbf{w} = k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + \dots + k_r\mathbf{v}_r$, where each k_i is a scalar. The scalars are called the **coefficients** of the linear combination. In the case where $r = 1$, we have $\mathbf{w} = k_1\mathbf{v}_1$, in which case the linear combination is a scalar multiple of the vector.

#2 Express the following as linear combinations of $\mathbf{u} = (2, 1, 4)$,
 $\mathbf{v} = (1, -1, 3)$, and $\mathbf{w} = (3, 2, 5)$.

a. $(-9, -7, -15)$

b. $(6, 11, 6)$

c. $(0, 0, 0)$

$$a \vec{u} + b \vec{v} + c \vec{w} = a \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix} + b \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} + c \begin{bmatrix} 3 \\ 2 \\ 5 \end{bmatrix} = \begin{bmatrix} -9 \\ -7 \\ -15 \end{bmatrix}$$

a) yields $\begin{bmatrix} 2 & 1 & 3 \\ 1 & -1 & 2 \\ 4 & 3 & 5 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} -9 \\ -7 \\ -15 \end{bmatrix}$

$\left[\begin{array}{ccc|c} 2 & 1 & 3 & -9 \\ 1 & -1 & 2 & -7 \\ 4 & 3 & 5 & -15 \end{array} \right]$ Faster than doing this repeatedly,

$$\left[\begin{array}{ccc|c} 2 & 1 & 3 & b_1 \\ 1 & -1 & 2 & b_2 \\ 4 & 3 & 5 & b_3 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 0 & -\frac{11}{2}b_1 + 2b_2 + \frac{5}{2}b_3 \\ 0 & 1 & 0 & \frac{3}{2}b_1 - b_2 - \frac{1}{2}b_3 \\ 0 & 0 & 1 & \frac{3}{2}b_1 - b_2 - \frac{3}{2}b_3 \end{array} \right]$$

Since there are no restrictions on b_1, b_2, b_3 , the vectors $\vec{u}, \vec{v}, \& \vec{w}$ span \mathbb{R}^3 .

In particular, if $b_1 = -9$, $b_2 = -7$, $b_3 = -15$,
then for $a\vec{u} + b\vec{v} + c\vec{w} = (-9, -7, -15)$,

$a = -2$, $b = 1$, $c = -2$ (we can check this)

Appropriate values for b_1, b_2, b_3 provide
solutions to parts (b) & (c).

Definition: The **span** of a nonempty set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ of vectors is the set of all possible linear combinations of vectors in S , denoted by $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ or $\text{span}(S)$. If $W = \text{span}(S)$, then we say that W is **spanned** by S .

#9 Determine whether the following polynomials span P_2 .

$$p_1 = 1 - x + 2x^2, p_2 = 3 + x, p_3 = 5 - x + 4x^2, p_4 = -2 - 2x + 2x^2$$

That is, can we build every polynomial of $\deg \leq 2$ using $\vec{p}_1, \vec{p}_2, \vec{p}_3, \& \vec{p}_4$?

Consider $\vec{p} = a_0 + a_1x + a_2x^2$.

Want $a\vec{p}_1 + b\vec{p}_2 + c\vec{p}_3 + d\vec{p}_4 = \vec{p}$.

$$\left[\begin{array}{cccc|c} 1 & 3 & 5 & -2 & a_0 \\ -1 & 1 & -1 & -2 & a_1 \\ 2 & 0 & 4 & 2 & a_2 \end{array} \right] \xrightarrow{\text{row reduce}} \left[\begin{array}{cccc|c} 1 & 0 & 2 & 1 & \frac{1}{4}a_0 - \frac{3}{4}a_1 \\ 0 & 1 & 1 & -1 & \frac{1}{4}a_0 + \frac{1}{4}a_1 \\ 0 & 0 & 0 & 0 & a_0 - 3a_1 - 2a_2 \end{array} \right]$$

R_3 restricts a_0, a_1, a_2

The set $\{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4\}$ does not

span P_2 .

#12 Let $T_A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be multiplication by A . Determine whether the vector $\mathbf{u} = (1, 2)$ is in the span of $\{T_A(\mathbf{e}_1), T_A(\mathbf{e}_2)\}$.

a. $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$

b. $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

a.

$$T(\vec{e}_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T(\vec{e}_2) = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$a \begin{bmatrix} 1 \\ 0 \end{bmatrix} + b \begin{bmatrix} 2 \\ 1 \end{bmatrix} \stackrel{?}{=} \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad b = 2, a = -3$$

yes

b. $a \begin{bmatrix} 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 1 \\ 1 \end{bmatrix} \stackrel{?}{=} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ no.

#16 Let W be the solution space to the system $A\mathbf{x} = \mathbf{0}$. Determine whether the set $\{\mathbf{u}, \mathbf{v}\}$ spans W .

$$A = \begin{bmatrix} 0 & 1 & -1 & 1 \\ 0 & 2 & -2 & 2 \\ 0 & 3 & -3 & 3 \end{bmatrix}$$

$$\xrightarrow{\text{ref}} \begin{bmatrix} 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \rightarrow x_2 = x_3 - x_4$$

leading

a. $\mathbf{u} = (1, 1, 1, 0), \mathbf{v} = (0, -1, 0, 1)$

b. $\mathbf{u} = (0, 1, 1, 0), \mathbf{v} = (1, 0, 1, 1)$

free: x_1, x_3, x_4

Let $r = x_1, s = x_3, t = x_4$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} r \\ s-t \\ s \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} r + \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} s + \begin{bmatrix} 0 \\ -1 \\ 0 \\ 1 \end{bmatrix} t$$

The solution space is span $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 0 \\ 1 \end{bmatrix} \right\}$

a) $\vec{u} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$, $\vec{v} = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$. No. We can't

get, for instance, $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ using \vec{u} & \vec{v} .

b) You'll find the answer is also no.

Theorem 4.3.1 If $S = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_r\}$ is a nonempty set of vectors in a vector space V , then:

a) The set W of all possible linear combinations of the vectors in S is a subspace of V .

b) The set W in part (a) is the "smallest" subspace of V that contains all of the vectors in S in the sense that any other subspace that contains those vectors contains W .

pf (a): Clearly $\vec{0} \in \text{span}(S)$. (nonempty)

Let $\vec{u}, \vec{v} \in \text{span}(S)$. Then $\vec{u} = \sum_{i=1}^r c_i \vec{w}_i$

and $\vec{v} = \sum_{i=1}^r k_i \vec{w}_i$. $\vec{u} + \vec{v} = \sum_{i=1}^r c_i \vec{w}_i + \sum_{i=1}^r k_i \vec{w}_i$

$= \sum_{i=1}^r (c_i + k_i) \vec{w}_i$, a linear comb.

$\Rightarrow \vec{u} + \vec{v} \in \text{span}(S)$.

$k\vec{u} = k \sum_{i=1}^r c_i \vec{w}_i = \sum_{i=1}^r (kc_i) \vec{w}_i \Rightarrow k\vec{u} \in \text{span}(S)$

$\text{span}(S)$ is a subspace.

Theorem 4.3.2 If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ and $S' = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_k\}$ are nonempty sets of vectors in a vector space V , then $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\} = \text{span}\{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_k\}$ if and only if each vector in S is a linear combination of those in S' , and each vector in S' is a linear combination of those in S .

#20 Let $\mathbf{v}_1 = (1, 6, 4)$, $\mathbf{v}_2 = (2, 4, -1)$, $\mathbf{v}_3 = (-1, 2, 5)$, and $\mathbf{w}_1 = (1, -2, -5)$, $\mathbf{w}_2 = (0, 8, 9)$. Use Theorem 4.3.2 to show that $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\} = \text{span}\{\mathbf{w}_1, \mathbf{w}_2\}$.

$$a\vec{v}_1 + b\vec{v}_2 + c\vec{v}_3 = \vec{w}_1$$

Can we use the \vec{v}_i to make all the \vec{w}_i

AND can we use the \vec{w}_i to make all the

\vec{v}_i ?

$$\begin{array}{c} \vec{v}_1 \quad \vec{v}_2 \quad \vec{v}_3 \quad \vec{w}_1 \quad \vec{w}_2 \\ \left[\begin{array}{ccc|cc} 1 & 2 & -1 & 1 & 0 \\ 6 & 4 & 2 & -2 & 8 \\ 4 & -1 & 5 & -5 & 9 \end{array} \right] \rightarrow \left[\begin{array}{ccc|cc} 1 & 0 & 1 & -1 & 2 \\ 0 & 1 & -1 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \end{array}$$

$$\vec{w}_1 = -\vec{v}_1 + \vec{v}_2 + 0\vec{v}_3 \quad \vec{w}_2 = 2\vec{v}_1 - \vec{v}_2$$

$$a\vec{v}_1 + b\vec{v}_2 + c\vec{v}_3 = \vec{w}_1$$

$$\begin{bmatrix} 1 & 2 & -1 \\ 6 & 4 & 2 \\ 4 & -1 & 5 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ -5 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & -1 & 1 \\ 6 & 4 & 2 & -2 \\ 4 & -1 & 5 & -5 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \rightarrow \begin{array}{l} a = -1 - c \\ b = 1 + c \end{array}$$

$$(-1 - c, 1 + c, c)$$

$$\vec{v}_1 - \vec{v}_2 = \vec{v}_3$$

From letting

$$c = 0.$$

What about building \vec{v}_i using \vec{w}_i ?

$$\left[\begin{array}{cc|cc} 1 & 0 & 1 & 2 & -1 \\ -2 & 8 & 6 & 4 & 2 \\ -5 & 9 & 4 & -1 & 5 \end{array} \right] \rightarrow \left[\begin{array}{cc|cc} 1 & 0 & 1 & 2 & -1 \\ 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

$$\Rightarrow \vec{v}_1 = \vec{w}_1 + \vec{w}_2$$

$$\Rightarrow \vec{v}_2 = 2\vec{w}_1 + \vec{w}_2$$

$$\Rightarrow \vec{v}_3 = -\vec{w}_1$$