

6.2 – Angle and Orthogonality in Inner Product Spaces

Definition: The angle θ between vectors \mathbf{u} and \mathbf{v} in a real inner product space V is $\theta = \cos^{-1} \left(\frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{u}\| \|\mathbf{v}\|} \right)$.

#5 Find the cosine of the angle between A and B with respect to the standard inner product on M_{22} .

$$A = \begin{bmatrix} 2 & 6 \\ 1 & -3 \end{bmatrix}, B = \begin{bmatrix} 3 & 2 \\ 1 & 0 \end{bmatrix}$$

Theorem 6.2.1 Cauchy-Schwarz Inequality (generalization of Theorem 3.2.4)
If \mathbf{u} and \mathbf{v} are vectors in a real inner product space V , then $|\langle \mathbf{u}, \mathbf{v} \rangle| \leq \|\mathbf{u}\| \|\mathbf{v}\|$.

Definition: Two vectors \mathbf{u} and \mathbf{v} in a real inner product space V are **orthogonal** if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$.

#10 Show that the vectors are orthogonal with respect to the standard inner product on P_2 .

$$\mathbf{p} = 2 - 3x + x^2, \mathbf{q} = 4 + 2x - 2x^2$$

Theorem 6.2.3 Generalized Theorem of Pythagoras

If \mathbf{u} and \mathbf{v} are orthogonal vectors in a real inner product space, then

$$\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2.$$

Example: Let $\mathbf{p} = p(x) = 1 - 2x^2$, $\mathbf{q} = q(x) = 4 - 2x + x^2$, $\mathbf{r} = r(x) = x + 2x^2$ and let P_2 have the standard inner product.

a. Compute the following: $\langle \mathbf{p}, \mathbf{q} \rangle$ and $\langle \mathbf{q}, \mathbf{r} \rangle$.

b. Now let P_2 have the inner product $\langle \mathbf{p}, \mathbf{q} \rangle = a_0b_0 + a_1b_1 + ka_2b_2$. Find k so that \mathbf{p} and \mathbf{q} are orthogonal.

c. Using this new inner product, compute $\langle \mathbf{q}, \mathbf{r} \rangle$.

#17 Do there exist scalars k and l such that the vectors $\mathbf{p}_1 = 2 + kx + 6x^2$, $\mathbf{p}_2 = l + 5x + 3x^2$, $\mathbf{p}_3 = 1 + 2x + 3x^2$ are mutually orthogonal with respect to the standard inner product on P_2 ?

Definition: (inner product space analog of Definition 2 in Section 4.9) If W is a subspace of a real inner product space V , then the set of all vectors in V that are orthogonal to every vector in W is called the **orthogonal complement** of W and is denoted by the symbol W^\perp (pronounced “ W perp”).

Theorem 6.2.4 (generalization of Theorem 4.9.6 (a) and (b))

If W is a subspace of a real inner product space V , then:

- a) W^\perp is a subspace of V .
- b) $W \cap W^\perp = \{\mathbf{0}\}$.

Theorem 6.2.5 (generalization of Theorem 4.9.6(c))

If W is a subspace of a real finite-dimensional inner product space V , then the orthogonal complement of W^\perp is W , that is, $(W^\perp)^\perp = W$.

#27 Find a basis for the orthogonal complement of the subspace of R^4 spanned by the vectors $\mathbf{v}_1 = (1, 4, 5, 2)$, $\mathbf{v}_2 = (2, 1, 3, 0)$, $\mathbf{v}_3 = (-1, 3, 2, 2)$.

